

Decomposition theorem on matchable distributive lattices*

Heping Zhang¹, Dewu Yang^{1,2} and Haiyuan Yao¹

¹School of Mathematics and Statistics, Lanzhou University, Lanzhou, Gansu 730000, P. R. China, zhanghp@lzu.edu.cn, hyyao@lzu.edu.cn.

²School of Mathematics and Statistics, Henan University of Science and Technology, Luoyang, Henan 471003, P. R. China, dewuyang0930@163.com

Abstract

A distributive lattice structure $\mathbf{M}(G)$ has been established on the set of perfect matchings of a plane bipartite graph G . We call a lattice *matchable distributive lattice* (simply MDL) if it is isomorphic to such a distributive lattice. It is natural to ask which lattices are MDLs. We show that if a plane bipartite graph G is elementary, then $\mathbf{M}(G)$ is irreducible. Based on this result, a decomposition theorem on MDLs is obtained: a finite distributive lattice $\mathbf{L}$ is an MDL if and only if each factor in any cartesian product decomposition of $\mathbf{L}$ is an MDL. Two types of MDLs are presented: $J(\mathbf{m} \times \mathbf{n})$ and $J(\mathbf{T})$, where $\mathbf{m} \times \mathbf{n}$ denotes the cartesian product between m -element chain and n -element chain, and $\mathbf{T}$ is a poset implied by any orientation of a tree.

Key words: Perfect matching, Plane bipartite graph, Z -transformation graph, Distributive lattice, Decomposition theorem.

AMS 2010 Subject Classifications: 05C70, 05C90, 06D05, 92E10.

1 Introduction

Perfect matching of graphs is significant for theoretical chemistry and theoretical physics. This graph-theoretical concept coincides with that of the Kekulé structure of organic molecules. The Kekulé structure count can be used to predict the stability of benzenoid hydrocarbons. The carbon-skeleton of a benzenoid hydrocarbon is a hexagonal system, i.e. 2-connected plane graph every interior face of which is a regular hexagon of side length

*This work was supported by NSFC (grant no. 10831001).

unit. Since 1980's there have been developed a combinatorial object, the Z -transformation graph (or resonance graph) [23, 24] on the set perfect matchings of a hexagonal system, late extended to a general plane bipartite graph [27, 33, 34, 35]; see a recent survey [29]. Randić [15, 16] showed that the leading eigenvalue of the resonance graphs has a quite satisfactory correlation with the resonance energy of benzenoid hydrocarbons.

A domino tiling of a polygon in the plane corresponds to a perfect matching of a related graph. In theoretical physics, a domino is seen as a dimer, a diatomic molecule (as the molecule of hydrogen), and each tiling is seen as a possible state of a solid or a fluid. In 2003 Fournier [5] reintroduced Z -transformation graph under name “perfect matching graph” in investigating domino tiling spaces of Saldanha et al. [19]. E. Rémila [17, 18] established the distributive lattice structure on the set of domino tilings of a polygon by using Thurston’s *height function*. In general, a distributive lattice on the set of perfect matchings of a plane bipartite graph was presented in terms of Z -transformation digraph and the unit decomposition of alternating cycle systems with respect to a perfect matching [10].

Let G be a finite and simple graph with vertex-set $V(G)$ and edge-set $E(G)$. A *perfect matching* or *1-factor* of G is a set of independent edges which saturate all vertices of G . Let $\mathcal{M}(G)$ denote the set of 1-factors of G . A plane bipartite graph G is *elementary* [11] if G is connected and every edge is contained in some 1-factor; further *weakly elementary* [34, 20] if every alternating cycle with respect to some 1-factor together with its interior form an elementary subgraph.

For a plane bipartite graph G , the Z -transformation graph $Z(G)$ is defined as a graph on $\mathcal{M}(G)$: $M, M' \in \mathcal{M}(G)$ are joined by an edge if and only if they differ only in one cycle that is the boundary of an inner face of G .

To give an acyclic orientation of $Z(G)$ [33], a proper 2-coloring (white-black) of bipartite graph G is specified. For $M \in \mathcal{M}(G)$, a cycle C is said to be *M -alternating* if the edges of C appear alternately in and off M ; further *proper (improper)* [32] if every edge of C belonging to M goes from white (black) end-vertex to black (white) end-vertex along the clockwise orientation of C . Now Z -transformation digraph $\vec{Z}(G)$ is the orientation of $Z(G)$: an edge M_1M_2 of $Z(G)$ is oriented from M_1 to M_2 if the symmetric difference $M_1 \oplus M_2$ form a proper M_1 - and improper M_2 -alternating cycle (the boundary of an inner face).

Since $\vec{Z}(G)$ has no directed cycles [33], it naturally implies a partial ordering on $\mathcal{M}(G)$. This poset is denoted by $\mathbf{M}(G)$. Then its Hasse diagram is isomorphic to $\vec{Z}(G)$. Lam and Zhang [10] showed that $\mathbf{M}(G)$ is a finite distributive lattice (FDL) if G is weakly elementary. Further the first author of the present paper showed [28] that $\mathbf{M}(G)$ is direct

sum of at least two distributive lattices if G is non-weakly elementary. By applying such a lattice structure, Zhang et al. showed [30] that every connected resonance graph of plane bipartite graphs is a median graph, and extended Klavžar et al.'s result [9] in the case of cata-condensed benzenoid systems.

In different ways, Propp [14] established a distributive lattice structure on the set of c -orientations of a plane bipartite graph G ; Pretzel [13] provided a new proof to Propp's result. Similar structures were also given on the set of reachable configurations of an edge firing game [12], α -orientations of a planar graph [4], and flows of a planar graph [7].

In this paper we propose a problem: which distributive lattices are isomorphic to distributive lattice $\mathbf{M}(G)$ on the set of 1-factors of a plane bipartite graphs G ? A lattice is called *matchable distributive lattice* (simply MDL) if it is isomorphic to such a distributive lattice $\mathbf{M}(G)$. Non-matchable distributive lattices exist. We show that if a plane bipartite graph G is elementary, then $\mathbf{M}(G)$ is irreducible. Based on this result, a decomposition theorem on MDL is obtained (Theorem 3.8): a finite distributive lattice $\mathbf{L}$ is an MDL if and only if each factor in any cartesian product decomposition of $\mathbf{L}$ is an MDL. Finally, we present two types of irreducible MDLs by applying the *fundamental theorem for finite distributive lattices* (FTFDL): $J(\mathbf{m} \times \mathbf{n})$ and $J(\mathbf{T})$, where $\mathbf{m} \times \mathbf{n}$ denotes the cartesian product between m -element chain and n -element chain, and $\mathbf{T}$ is a poset implied by any orientation of a tree. Meantime, we also show that for any order ideal W of $\mathbf{m} \times \mathbf{n}$, $J(W)$ is an MDL.

2 Preliminaries

Terms on poset and distributive lattice used in this paper can be found in [1, 6, 21]. If $\mathbf{P}$ and $\mathbf{Q}$ are posets, then the direct (cartesian) product of $\mathbf{P}$ and $\mathbf{Q}$ is the poset $\mathbf{P} \times \mathbf{Q}$ on the set $\{(x, y) : x \in \mathbf{P} \text{ and } y \in \mathbf{Q}\}$ such that $(x, y) \preceq (x', y')$ in $\mathbf{P} \times \mathbf{Q}$ if $x \preceq x'$ in $\mathbf{P}$ and $y \preceq y'$ in $\mathbf{Q}$.

Let $\mathbf{L}$ be an FDL with the greatest element $\hat{1}$ and the least element $\hat{0}$. If $\mathbf{L}$ can be expressed as the direct product of a series of FDLs $\mathbf{L}_j (j \in J)$, i.e. $\mathbf{L} = \prod_{j \in J} \mathbf{L}_j$, then we say that $\mathbf{L}$ has a (*direct product*) *decomposition* $\prod_{j \in J} \mathbf{L}_j$. A lattice with exactly one element is viewed as a trivial lattice. An FDL is *irreducible* if it cannot be expressed as direct product of at least two non-trivial FDLs. A decomposition $\mathbf{L} = \prod_{j \in J} \mathbf{L}_j$ is called *irreducible* if each $\mathbf{L}_j (j \in J)$ is non-trivial and irreducible.

For a decomposition $\mathbf{L} = \prod_{i=1}^n \mathbf{L}_i$, let $\hat{1}_i$ and $\hat{0}_i$ denote the greatest element and the least element of $\mathbf{L}_i$, respectively. Then $\hat{0} = (\hat{0}_1, \hat{0}_2, \dots, \hat{0}_n)$ and $\hat{1} = (\hat{1}_1, \hat{1}_2, \dots, \hat{1}_n)$. If each $\mathbf{L}_i$




Figure 1. A pair of central elements $(2, 1)$ and $(1, 3)$ of direct product $\mathbf{2} \times \mathbf{3}$.

is non-trivial and $n \geq 2$, $e_i = (\hat{0}_1, \dots, \hat{0}_{i-1}, \hat{1}_i, \hat{0}_{i+1}, \dots, \hat{0}_n)$ is called a *central element* of $\mathbf{L}$. For $x, y \in \mathbf{L}$, x is called a complement of y if $x \vee y = \hat{1}$ and $x \wedge y = \hat{0}$. The complement of x , when it exists, is unique. For example, two central elements $(2, 1)$ and $(1, 3)$ of $\mathbf{L} = \mathbf{2} \times \mathbf{3}$ are complementary each other (see Fig. 1). For a positive integer n , $\{1, 2, \dots, n\}$ with its usual order forms an n -element chain, denoted by $\mathbf{n}$.

Lemma 2.1. [1] *Any central element of an FDL has a unique complement.*

Lemma 2.2. [1] *Any FDL has a unique irreducible decomposition, i.e. if both $\prod_{i=1}^n \mathbf{L}_i$ and $\prod_{j=1}^m \mathbf{L}'_j$ are irreducible decompositions of $\mathbf{L}$, then $m = n$ and there exists a permutation π of $[n]$ such that $\mathbf{L}_i = \mathbf{L}'_{\pi(i)}$ ($i = 1, 2, \dots, n$).*

For an FDL $\mathbf{L}$, its rank function [21] satisfies

$$\rho(x) + \rho(y) = \rho(x \wedge y) + \rho(x \vee y),$$

for any $x, y \in \mathbf{L}$. For a pair of complementary elements x and y of $\mathbf{L}$, we have

$$\rho(x) + \rho(y) = \rho(\hat{0}) + \rho(\hat{1}) = \rho(\hat{1}) = \rho(\mathbf{L}).$$

Lemma 2.3. *Let $\mathbf{L}$ be an FDL of rank k and y the complement of $x \in \mathbf{L}$. If $\rho(x) = r \geq 1$ and $\rho(y) = k - r \geq 1$, then $\mathbf{L}$ has a sublattice $(\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$ containing x and y .*

Proof. $\mathbf{L}$ has at least two saturated chains between $\hat{0}$ with x and y , respectively:

$$P_1 : \hat{0} = x_0 \prec x_1 \prec x_2 \prec \dots \prec x_r = x,$$

and

$$P_2 : \hat{0} = y_0 \prec y_1 \prec y_2 \prec \dots \prec y_{k-r} = y.$$

Then $x_i \wedge y_j = \hat{0}$, for any $0 \leq i \leq r, 0 \leq j \leq k - r$, since $x_i \wedge y_j \preceq x \wedge y = \hat{0}$. Hence P_1 and P_2 have no common elements except for $\hat{0}$.

Let $\mathbf{L}' = \{a_{ij} : a_{ij} = x_i \vee y_j, 0 \leq i \leq r, 0 \leq j \leq k - r\}$. Then $\mathbf{L}'$ satisfies the following three properties:

1. $a_{ij} = a_{i'j'}$ if and only if $i = i'$ and $j = j'$. If $a_{i'j'} \preceq a_{ij}$, i.e. $x_{i'} \vee y_{j'} \preceq x_i \vee y_j$, then $(x_{i'} \vee y_{j'}) \wedge y_{j'} \preceq (x_i \vee y_j) \wedge y_{j'}$. By distributive laws and $x_i \wedge y_j = \hat{0}$, we have that $y_{j'} \preceq y_j \wedge y_{j'}$ and $j' \leq j$. Similarly we have $i' \leq i$. So the property holds.

2. $\mathbf{L}'$ forms a sublattice of $\mathbf{L}$ and $\mathbf{L}' = \langle x_1, \dots, x_r, y_1, \dots, y_{k-r}; \vee, \wedge \rangle$. It suffices to show that $\mathbf{L}'$ is closed under meet and join operations $\wedge$ and $\vee$ of $\mathbf{L}$.

(i) $a_{ij} \vee a_{i'j'} = (x_i \vee y_j) \vee (x_{i'} \vee y_{j'}) = (x_i \vee x_{i'}) \vee (y_j \vee y_{j'}) = x_{i''} \vee y_{j''} = a_{i''j''} \in \mathbf{L}'$, where $i'' = \max\{i, i'\}$, $j'' = \max\{j, j'\}$;

(ii)

$$\begin{aligned} a_{ij} \wedge a_{i'j'} &= a_{ij} \wedge (x_{i'} \vee y_{j'}) = (a_{ij} \wedge x_{i'}) \vee (a_{ij} \wedge y_{j'}) \\ &= ((x_i \vee y_j) \wedge x_{i'}) \vee ((x_i \vee y_j) \wedge y_{j'}) \\ &= ((x_i \wedge x_{i'}) \vee (y_j \wedge x_{i'})) \vee ((x_i \wedge y_{j'}) \vee (y_j \wedge y_{j'})) \\ &= ((x_i \wedge x_{i'}) \vee \hat{0}) \vee (\hat{0} \vee (y_j \wedge y_{j'})) \\ &= x_{i'''} \vee y_{j'''} = a_{i'''j'''} \in \mathbf{L}', \end{aligned}$$

where $i''' = \min\{i, i'\}$, $j''' = \min\{j, j'\}$.

3. $\mathbf{L}'$ is isomorphic to $(\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$. Let $\phi : \mathbf{L}' \rightarrow (\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$ be a bijection as $\phi(a_{ij}) = (i, j)$ for any $a_{ij} \in \mathbf{L}'$. Then by Property 2(i) and (ii), we have that

$$\phi(a_{ij} \vee a_{i'j'}) = \phi(a_{i''j''}) = (i'', j'') = (i, j) \vee (i', j') = \phi(a_{ij}) \vee \phi(a_{i'j'}),$$

and

$$\phi(a_{ij} \wedge a_{i'j'}) = \phi(a_{i'''j'''}) = (i''', j''') = (i, j) \wedge (i', j') = \phi(a_{ij}) \wedge \phi(a_{i'j'}).$$

Hence $\mathbf{L}' \cong (\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$. □

3 Some fundamental results on MDL

Let G be a plane bipartite graph with a specific proper black-white coloring to vertices. An edge e of a cycle (or an inner face) C is *proper* if e goes from the white end-vertex to the black endvertex along the clockwise direction of C . Let $\mathcal{F}(G)$ denote the set of all inner faces of G . Recall that $\mathcal{M}(G)$ denotes the set of all 1-factors of G .

Definition 1. A binary relation $\preceq$ on $\mathcal{M}(G)$ is defined as: $M_1 \preceq M_2$, $M_1, M_2 \in \mathcal{M}(G)$, if and only if $\vec{Z}(G)$ has a directed path from M_2 to M_1 .

It is known that $\mathbf{M}(G) = (\mathcal{M}(G), \preceq)$ is a poset and a lattice structure on $\mathcal{M}(G)$ is revealed in the following two theorems.

Theorem 3.1. [10] *Let G be a plane (weakly) elementary bipartite graph. Then $\mathbf{M}(G)$ is a finite distributive lattice, and its Hasse diagram is isomorphic to $\vec{Z}(G)$.*

Theorem 3.2. [28] *Let G be a plane bipartite graph with 1-factor. Then $\mathbf{M}(G)$ is direct sum of distributive lattices and the Hasse diagram is isomorphic to $\vec{Z}(G)$.*

Definition 2. An FDL $\mathbf{L}$ is called an *matchable distributive lattice* (MDL) if there exist a plane bipartite graph G such that $\mathbf{L} \cong \mathbf{M}(G)$.

Let $M^{\hat{1}}$ and $M^{\hat{0}}$ denote 1-factors of G such that G has neither improper $M^{\hat{1}}$ - nor proper $M^{\hat{0}}$ -alternating cycles, called *source* and *root* 1-factors of G respectively. If $\mathbf{M}(G)$ is an FDL, then $M^{\hat{1}}$ and $M^{\hat{0}}$ are the greatest element and the least element, respectively.

Lemma 3.3. *Let G be a plane elementary bipartite graph with more than two vertices. Then the boundary of G is proper $M^{\hat{1}}$ - and improper $M^{\hat{0}}$ -alternating cycle.*

Proof. It is known that G is 2-connected and the boundary is a cycle. For every proper edge $e = uv$ on the boundary of G , it suffices to show that $e \in M^{\hat{1}}$. Otherwise, an edge e' different from e and incident to u belongs to $M^{\hat{1}}$. Since G is elementary, it has a 1-factor M such that $e \in M$. Then $M \oplus M^{\hat{1}}$ has a cycle containing e and e' , which is both improper $M^{\hat{1}}$ - and proper M -alternating cycle, a contradiction. Hence the boundary of G is proper $M^{\hat{1}}$ -alternating cycle. Similarly, we can show that the boundary of G is improper $M^{\hat{0}}$ -alternating cycle. $\square$

Let G be a plane elementary bipartite graph with $M' \preceq M$ in $\mathbf{M}(G)$. For any $f \in \mathcal{F}(G)$, let $\Delta_{\mathcal{C}}(f)$ denote the number of proper M -alternating cycles in $\mathcal{C} := \mathcal{C}(M, M') = M \oplus M'$ with f in their interiors minus the number of improper M -alternating cycles in $\mathcal{C}$ with f in their interiors. Then $M' \preceq M$ implies that $\Delta_{\mathcal{C}}(f) \geq 0$ by Lemma 3.1 in [28]. For any directed path $\vec{P} = M_0(= M)M_1 \dots M_t(= M')$ from M to M' of $\vec{Z}(G)$, let $s_i := M_{i-1} \oplus M_i$, $i = 1, \dots, t-1$. Let $\delta_P(f)$ denote the times of f appearing in the face sequence corresponding to $s_1, \dots, s_t$. Lemma 3.5 in Ref. [28] implies the following result.

Lemma 3.4. *Let G be a plane elementary bipartite graph with $M' \preceq M$ in $\mathbf{M}(G)$. If $\vec{P}$ is a directed path from M to M' of $\vec{Z}(G)$ and $\mathcal{C} = M \oplus M'$, then $\delta_P(f) = \Delta_{\mathcal{C}}(f)$ for each $f \in \mathcal{F}(G)$.*

From Lemmas 2.3 and 3.3 we can derive the following critical result.

Lemma 3.5. *For a plane elementary bipartite graph G with more than two vertices, each element of $\mathbf{M}(G)$ has no complement except the greatest element $M^{\hat{1}}$ and the least element $M^{\hat{0}}$.*




Figure 2. Sublattice $(\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$.

Proof. Suppose to the contrary that $\mathbf{M}(G)$ has a pair of mutually complementary elements M and M' except $M^{\hat{1}}$ and $M^{\hat{0}}$. Let $\rho(\mathbf{M}(G)) = k$ and $\rho(M) = r$. Then $\rho(M') = k - r$, and $k - 1 \geq r \geq 1$. By Lemma 2.3, $\mathbf{M}(G)$ has a sublattice $(\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$ as shown in Fig. 2 containing the following two maximal chains:

$$M_0(= M^{\hat{0}}) \prec M_1 \prec \cdots \prec M_r(= M) \prec \cdots \prec M_{k-1} \prec M_k(= M^{\hat{1}}), \text{ and}$$

$$M'_0(= M^{\hat{0}}) \prec M'_1 \prec \cdots \prec M'_{k-r}(= M') \prec \cdots \prec M'_{k-1} \prec M'_k(= M^{\hat{1}}).$$

Put $M_{ij} := M_i \vee M'_j$, $i = 0, 1, \dots, r$, $j = 0, 1, \dots, k - r$, $s_i := M_i \oplus M_{i-1}$ ($1 \leq i \leq r$) and $s'_j := M'_j \oplus M'_{j-1}$ ($1 \leq j \leq k - r$). Since each maximal chain of $(\mathbf{r} + \mathbf{1}) \times (\mathbf{k} - \mathbf{r} + \mathbf{1})$ is a saturated chain of $\mathbf{M}(G)$, the s_i and s'_j are the boundaries of inner faces of G .

Claim 1. $M_{i,j} = M_{i-1,j} \oplus s_i = M_{i,j-1} \oplus s'_j$, and s_i and s'_j are disjoint, for $i = 1, 2, \dots, r$, and $j = 1, 2, \dots, k - r$.

Proof. We prove that $M_{i,j} = M_{i-1,j} \oplus s_i = M_{i,j-1} \oplus s'_j$ such that s_i and s'_j are proper $M_{i,j}$ -alternating by induction on $(i, j) \geq (1, 1)$. For $i = j = 1$, s_1 and s'_1 are improper $M^{\hat{0}}$ -alternating facial cycles, and are thus disjoint. Hence we have that $M_{1,1} = M_1 \oplus s'_1 = M'_1 \oplus s_1$

by Lemma 2.3, and the required holds. Let $i \geq 2$ or $j \geq 2$. For the induction step, suppose that the assertion holds for smaller i or j . By induction hypothesis, $M_{i,j-1} = M_{i-1,j-1} \oplus s_i$ and $M_{i-1,j} = M_{i-1,j-1} \oplus s'_j$, and s_i and s'_j are distinct and improper $M_{i-1,j-1}$ -alternating facial cycles, and are disjoint. Hence s_i and s'_j are improper $M_{i,j-1}$ -alternating and improper $M_{i-1,j}$ -alternating, respectively, and $M_{i,j-1} \oplus s'_j$ and $M_{i-1,j} \oplus s_i$ cover $M_{i,j-1}$ and $M_{i-1,j}$, respectively. Obviously, $M_{i,j-1} \oplus s'_j = M_{i-1,j-1} \oplus s_i \oplus s'_j = M_{i-1,j-1} \oplus s'_j \oplus s_i = M_{i-1,j} \oplus s_i$. Hence $M_{i,j} = M_{i-1,j} \oplus s_i = M_{i,j-1} \oplus s'_j$ since $M_{i,j} = M_{i-1,j} \vee M_{i,j-1}$ by Lemma 2.3. The assertion holds for any (i, j) . $\square$

Claim 2. Let f_i and h_j denote the inner faces of G bounded by s_i and s'_j , respectively. Then $\mathcal{F}(G) = \{f_1, f_2, \dots, f_r, h_1, h_2, \dots, h_{k-r}\}$.

Proof. Let $\mathcal{F}_1 := \{f_1, f_2, \dots, f_r\}$ and $\mathcal{F}_2 := \{h_1, h_2, \dots, h_{k-r}\}$. So we want to prove that $\mathcal{F}(G) = \mathcal{F}_1 \cup \mathcal{F}_2$.

Let $\mathcal{C} := M^{\hat{1}} \oplus M^{\hat{0}}$. Then each cycle in $\mathcal{C}$ is proper $M^{\hat{1}}$ and improper $M^{\hat{0}}$ -alternating cycle, one being the boundary of G by Lemma 3.3. Hence $\Delta_{\mathcal{C}}(f) \geq 1$ for any $f \in \mathcal{F}$.

Let $P := M_{r,k-r}(= M^{\hat{1}})M_{r,k-r-1} \cdots M_{r,0}M_{r-1,0} \cdots M_{0,0}(= M^{\hat{0}})$ be a directed path of $\vec{Z}(G)$, corresponding to a maximal chain of $\mathbf{M}(G)$. For any $f \in \mathcal{F}$, by Lemma 3.4 we have that $\delta_P(f) = \Delta_{\mathcal{C}}(f) \geq 1$. Hence $\mathcal{F}(G) = \{f_1, f_2, \dots, f_r, h_1, h_2, \dots, h_{k-r}\}$. $\square$

Since G is 2-connected, inner dual graph $G^{\#}$ of G is connected. Let f^* be a vertex of $G^{\#}$ corresponding to $f \in \mathcal{F}$. Then there must exist a vertex f_i^* in $\{f_1^*, \dots, f_r^*\}$ being adjacent to a vertex h_j^* in $V(G^*) \setminus \{f_1^*, \dots, f_r^*\} = \{h_1^*, \dots, h_{k-r}^*\}$. That means that f_i and h_j are adjacent, contradicting Claim 1. $\square$

From the above arguments, we have the following main results of this paper.

Theorem 3.6. *For a plane elementary bipartite graph G , $\mathbf{M}(G)$ is irreducible.*

Proof. If $G = K_2$, it is trivial. Otherwise, $\mathbf{M}(G)$ is a non-trivial FDL. By Lemma 3.5, every element of $\mathbf{M}(G)$ has no complement except for $M^{\hat{1}}$ and $M^{\hat{0}}$. By Lemma 2.1, $\mathbf{M}(G)$ has no central elements. Hence, $\mathbf{M}(G)$ is irreducible. $\square$

Elementary components of a plane bipartite graph G with 1-factor mean components other than K_2 of the subgraph obtained from G by the removal of all forbidden edges (those edges not contained in any 1-factors).

Corollary 3.7. *Let G be a weakly elementary plane bipartite graph with elementary components $G_1, G_2, \dots, G_k$. Then $\mathbf{M}(G) = \mathbf{M}(G_1) \times \mathbf{M}(G_2) \times \cdots \times \mathbf{M}(G_k)$ is an irreducible decomposition.*

Theorem 3.8. (*Decomposition Theorem*) Let $\mathbf{L}$ be an FDL with a decomposition $\mathbf{L} = \prod_{i=1}^n \mathbf{L}_i$. Then $\mathbf{L}$ is an MDL if and only if each $\mathbf{L}_i (1 \leq i \leq n)$ is an MDL.

Proof. If each factor $\mathbf{L}_i$ is an MDL, $1 \leq i \leq n$, then there exists a weakly elementary plane bipartite graph G_i such that $\mathbf{M}(G_i) \cong \mathbf{L}_i$. We construct a weakly elementary plane bipartite graph G by connecting G_i to G_{i+1} with a new edge in their exteriors for each $1 \leq i \leq n-1$. Then such new edges are forbidden edges of G . It follows that $\mathbf{M}(G) \cong \mathbf{M}(G_1) \times \mathbf{M}(G_2) \times \cdots \times \mathbf{M}(G_n) \cong \mathbf{L}_1 \times \mathbf{L}_2 \times \cdots \times \mathbf{L}_n = \mathbf{L}$. Hence $\mathbf{L}$ is an MDL.

Conversely, suppose that $\mathbf{L}$ is an MDL. Then there exists a plane weakly elementary bipartite graph G such that $\mathbf{M}(G) \cong \mathbf{L}$. Let $G_1, \dots, G_m$ be the non-trivial elementary components of G ($m \geq 1$). By Corollary 3.7 $\mathbf{L} \cong \prod_{j=1}^m \mathbf{M}(G_j)$ is an irreducible decomposition. If $\mathbf{L} = \prod_{i=1}^n \mathbf{L}_i$ is irreducible, then by Lemma 2.2, $m = n$ and there exists a permutation π of $[n]$ such that $\mathbf{L}_i = \mathbf{M}(G_{\pi(i)})$ $i = 1, 2, \dots, n$. So each $\mathbf{L}_i$ ($1 \leq i \leq n$) is an MDL. If $\prod_{i=1}^n \mathbf{L}_i$ is not irreducible, then each factor $\mathbf{L}_i$ is a direct product of some $\mathbf{M}(G_j)$'s. So each factor $\mathbf{L}_i$ is still an MDL. $\square$

4 MDL $\mathbf{J}(\mathbf{m} \times \mathbf{n})$

From now on we will present two typical irreducible MDLs by the *fundamental theorem for finite distributive lattice* (FTFDL).

Let $\mathbf{P}$ be a finite poset. An *order ideal* (*semi-ideal* or *down-set*) $\mathbf{I}$ of $\mathbf{P}$ is a subset of $\mathbf{P}$ if for every $x \in \mathbf{I}$, $y \preceq x$ implies $y \in \mathbf{I}$. The set $J(\mathbf{P})$ of order ideals of $\mathbf{P}$, ordered by the set-inclusion, forms a poset $\mathbf{J}(\mathbf{P})$. It is well known that $\mathbf{J}(\mathbf{P})$ is indeed a distributive lattice. The FTFDL states that the converse is true.

Theorem 4.1 (FTFDL). ([21]) Let $\mathbf{L}$ be an FDL. Then there is a unique (up to isomorphism) finite poset $\mathbf{P}$ for which $\mathbf{L} \cong \mathbf{J}(\mathbf{P})$.

In fact the above $\mathbf{P}$ can be viewed as a subposet of $\mathbf{L}$ consisting of all join-irreducible elements of $\mathbf{L}$: an element x of $\mathbf{L}$ is said to be *join-irreducible* if one cannot write $x = y \vee z$ where $y \prec x$ and $z \prec x$.

In this section we show that $\mathbf{J}(W)$ are MDLs for any order ideal W of $\mathbf{m} \times \mathbf{n}$. Let us introduce a type of hexagonal systems called truncated parallelogram [2, 3]: A *truncated parallelogram*, simply denoted by $H := L(r_1, r_2, \dots, r_m)$, consists of m condensed linear chains (rows) of the length $r_1, \dots, r_m$, $r_1 \geq r_2 \geq \dots \geq r_m > 0$ and the first hexagons (conventionally drawn to the left) from all chains also form a linear chain, the first column;




Figure 3. (a) Truncated parallelogram $H = L(r_1, r_2, \dots, r_m)$ with root 1-factor $M^{\hat{0}}$, (b) parallelogram $L(m; n)$, and (c) prolate triangle T_m .

In particular, $L(m; n) = L(n, n, \dots, n)$ is a *parallelogram*, and $T_m := L(m, m-1, \dots, 1)$ is a *prolate triangle*. For example, see Fig. 3. For convenience, all hexagonal systems considered in this section are drawn such that an edge-direction is vertical and the valleys are colored white.

Let L and B be the left and bottom perimeters of H , respectively, which have a black vertex in common. The root 1-factor $M^{\hat{0}}$ of H has all vertical edges in L , and a series of parallel edges of B from left-low to right-up, and a series of parallel edges of $H - L - B$ from left-up to right-lower. We can see that the boundary of H is an improper $M^{\hat{0}}$ -alternating cycle. Hence H is elementary [34].

Since H has a forcing edge e (an edge contained in a unique 1-factor), each $M^{\hat{0}}$ -alternating cycle must pass through e ; see [25] for details. For each 1-factor M of H other than $M^{\hat{0}}$, $C_M := M \oplus M^{\hat{0}}$ is an $M^{\hat{0}}$ -alternating cycle of H . Thus we have a bijection [26] between the 1-factors other than $M^{\hat{0}}$ of H and the $M^{\hat{0}}$ -alternating cycles of H . Hence the subhexagonal system of H formed by C_M together with its interior is also a truncated parallelogram. Conversely, the perimeter of any sub-truncated parallelogram of H with edge e is an $M^{\hat{0}}$ -alternating cycle. Hence each 1-factor M of H corresponds exactly to a sub-truncated parallelogram of H with edge e , denoted by H_M . However, $H_{M^{\hat{0}}}$ corresponds to the empty graph (without vertex), the degenerated sub-truncated parallelogram of H .

Let $P_M := (L \cup B) \oplus C_M$. Then P_M is an M -alternating path with both end-edges in M (see Fig. 4(a)). Note that $C_{M^{\hat{0}}} = \emptyset$ and $P_{M^{\hat{0}}} = L \cup B$. From $M = M^{\hat{0}} \oplus C_M$, we have the following structure of M .

Proposition 4.2. *For each $M \in \mathcal{M}(H)$, the edges in $M \setminus E(P_M)$ have the same edge-direction from left-up to right-low.* $\square$




Figure 4. (a) Truncated parallelogram H with an $M, M^{\hat{0}}$ -alternating cycle C_M , (b) M -alternating path P_M , (c) a hexagon h , and (d) sub-parallelogram $H_{M'}$.

Lemma 4.3. *Let $M \in \mathcal{M}(H)$ and h an M -alternating hexagon of H . Then h intersects at three consecutive edges of P_M ; Moreover, h is proper if and only if $h \subseteq H_M$.*

Proof. If h is disjoint with P_M , then h is not M -alternating by Proposition 4.2. Otherwise, $1 \leq |E(h \cap P_M)| \leq 3$. Since h is M -alternating, h intersects at three consecutive edges of P_M . So h is proper M -alternating if and only if $e_2, e_4, e_6 \in M$. This holds if and only if $e_2, e_6 \in M \cap E(P_M)$. Thus h and P_M have exactly three common edges e_1, e_2, e_6 and $h \in H_M$. Similarly, h is improper M -alternating if and only if h and P_M have exactly three common edges e_3, e_4, e_5 and $h \notin H_M$ (see Fig. 4 (b) and (c)). $\square$

Lemma 4.4. *Let $M, M' \in \mathcal{M}(H)$. Then $M' \preceq M$ in $\mathbf{M}(H)$ if and only if $H_{M'}$ is a sub-truncated parallelogram of H_M , namely $H_{M'} \subseteq H_M$.*

Proof. We first show that M covers M' if and only if H_M can be obtained from $H_{M'}$ by adding a hexagon. If M covers M' , then there is a proper M -alternating hexagon h such that $M' = M \oplus h$ by Theorem 3.1, and $C_{M'} = M' \oplus M^{\hat{0}} = (M' \oplus M) \oplus (M \oplus M^{\hat{0}}) = h \oplus C_M$. By Lemma 4.3, we have that $h \in H_M$ has exactly three edges of P_M . If $h = C_M$, the result is trivial. Otherwise, $C_{M'}$ is an improper $M^{\hat{0}}$ -alternating cycle, and the sub-truncated parallelogram $H_{M'}$ of H bounded by $C_{M'}$ can be obtained by removing h from H_M .

Conversely, assume that $H_{M'}$ can be obtained from H_M by removing a hexagon h of H . Since both H_M and $H_{M'}$ are sub-truncated parallelograms of H , $h \in H_M$ must have exactly three edges of P_M . By Lemma 4.3, h is proper M -alternating. Then $M' = M^{\hat{0}} \oplus C_{M'} = M^{\hat{0}} \oplus (C_M \oplus h) = M \oplus h$. Hence M covers M' in $\mathbf{M}(H)$.

We now show the lemma. If $M' \preceq M$ in $\mathbf{M}(H)$, we can show that $H_{M'} \subseteq H_M$ by choosing a saturated chain between M and M' and applying repeatedly the above fact proved. If $H_{M'} \subseteq H_M$, there are a series of sub-truncated parallelograms of H : $H_1 (= H_{M'}), H_2, \dots, H_t (= H_M)$, such that each H_i is obtained from H_{i+1} by adding a hexagon.

Each H_i corresponds to a 1-factor M_i of H , $i = 1, 2, \dots, t$. By the above fact we have M_{i+1} covers M_i , $i = 1, 2, \dots, t-1$. Hence $M' \preceq M$. $\square$

Now, we define a poset on $\mathcal{F}(H)$, the set of hexagons of H . $h \in H$ is labeled with h_{ij} if h lies in the i -th row and j -th column, $1 \leq j \leq r_i, 1 \leq i \leq m$. For two hexagons h_{ij} and h_{kl} , $h_{ij} \preceq h_{kl}$ if and only if $i \leq k$ and $j \leq l$. Then $\mathbf{F}(H) := (\mathcal{F}(H), \preceq)$ is a poset. If H is a parallelogram, then $\mathbf{F}(H)$ is $\mathbf{m} \times \mathbf{n}$. In general, $\mathbf{F}(H)$ is an order ideal of $\mathbf{m} \times \mathbf{n}$. For example, see Fig. 5.




Figure 5. (a) $\mathbf{F}(L(r_1, r_2, \dots, r_m))$, (b) $\mathbf{F}(L(m; n))$, and (c) $\mathbf{F}(T_m)$.

By Lemma 4.4, we can see that $M \in \mathbf{M}(H)$ is join-irreducible if and only if H has a unique proper M -alternating hexagon lying in H_M , which is a sub-parallelogram of H . Let $\mathcal{I}(\mathbf{M}(H))$ denote the subposet of $\mathbf{M}(G)$ consisting of all join-irreducible elements.

Lemma 4.5. $\mathcal{I}(\mathbf{M}(H)) \cong \mathbf{F}(H)$.

Proof. A bijection $\psi : \mathcal{I}(\mathbf{M}(H)) \rightarrow \mathbf{F}(H)$ is defined as follows. For each $M \in \mathcal{I}(\mathbf{M}(H))$, let $\psi(M)$ denote the unique proper M -alternating hexagon of H_M , i.e. the right-up-most hexagon of H_M . Moreover, both ψ is an isomorphism: for any $M, M' \in \mathcal{I}(\mathbf{M}(H))$, by Lemma 4.4 we have that $M' \preceq M$ in $\mathbf{M}(H) \Leftrightarrow H_{M'} \subseteq H_M \Leftrightarrow \psi(M') \preceq \psi(M)$ in $\mathbf{F}(H)$. $\square$

By Theorem 4.1 and Lemma 4.5, we have a main theorem as follows.

Theorem 4.6. $\mathbf{M}(H) \cong \mathbf{J}(\mathbf{F}(H))$. $\square$

When H takes all over the truncated parallelograms for fixed m and n , $\mathbf{F}(H)$ goes all order ideals of $\mathbf{m} \times \mathbf{n}$. From the above theorem we have an immediate consequence as follows.

Corollary 4.7. *Let W be any order ideal of $\mathbf{m} \times \mathbf{n}$. Then $\mathbf{J}(W)$ is an irreducible MDL. $\square$*

We can obtain a series of MDLs by applying the above theorem to the special truncated parallelograms, such as parallelogram, prolate triangle, etc. Let $\mathbf{R}_m := \mathbf{F}(T_m)$. Here we give two special ones:

Corollary 4.8. *$\mathbf{J}(\mathbf{m} \times \mathbf{n})$ and $\mathbf{J}(\mathbf{R}_m)$ are irreducible MDLs. Moreover*

$$(1) \mathbf{J}(\mathbf{m} \times \mathbf{n}) \cong \mathbf{M}(L(m; n)), \text{ and}$$

$$(2) \mathbf{J}(\mathbf{R}_m) \cong \mathbf{M}(T_m). \quad \square$$

Note that [2, 3] the number of 1-factors of parallelogram $L(m; n)$ and prolate triangle T_m are $\binom{m+n}{m}$ and $\frac{1}{m+2} \binom{2m+2}{m+1}$ (Catalan number), respectively.

5 MDL $\mathbf{J}(\mathbf{T})$

In this section we will show that $\mathbf{J}(\mathbf{T})$ is an irreducible MDL with outerplane bipartite graphs for a poset $\mathbf{T}$ implied by any orientation of a tree. A connected plane graph G is *outerplane* if all vertices lie on the boundary of the outer face of G . Let $\mathcal{G}$ be the set of all 2-connected outerplane bipartite graphs. Catacondensed hexagonal systems are typical members of $\mathcal{G}$ [8].

An edge set T of a connected graph G is called a *minimal* edge-cut if $G - T$ is not connected but $G - T'$ remains connected for any proper subset T' of T . For a plane graph G , let e^* and f^* denote the edge and vertex of dual graph G^* corresponding to edge e and face f of G , respectively; For $T \subseteq E(G)$, put $T^* := \{e^* : e \in T\}$. Some edges in a plane graph G form a minimal edge-cut in G if and only if the corresponding dual edges form a cycle in G^* [22]. A minimal edge-cut T of a plane bipartite graph G is called *elementary edge cut* (*e-cut* for short) [34] if all edges of T are incident with white vertices of one component of $G - T$, called the *white bank* of T , and the other component is the *black bank* of T .

Lemma 5.1. [31] *Let T be a minimal edge-cut of $G \in \mathcal{G}$. Then T is an e-cut of G if and only if for any 1-factor M of G , $|M \cap T| = 1$.*

We now give an orientation $\vec{G}^*$ of the dual G^* : an edge e^* is oriented as an arc from f_1^* to f_2^* if one goes along e^* from f_1^* to f_2^* the white end-vertex of e lies right side. For example, see Figs. 6 and 7. We can see that a minimal edge-cut T is an e-cut of G if and only if T^* forms a directed cycle of $\vec{G}^*$.

For $G \in \mathcal{G}$, we now give a poset on $\mathcal{F}(G)$. Let $\vec{G}^\#$ be the orientation of inner dual graph $G^\#$, obtained from directed dual graph $\vec{G}^*$ by deleting the vertex f_0^* corresponding to the outer face of G . For $f_1, f_2 \in \mathcal{F}(G)$, we define " $f_1 \leq f_2$ " if $\vec{G}^\#$ has a directed path from f_2^* to f_1^* . Since $\vec{G}^\#$ contains no directed cycles, $\mathbf{F}(G) := (\mathcal{F}(G), \leq)$ is a poset.

For a plane elementary bipartite graph G with $M, M' \in \mathbf{M}(G)$, if $M' \preceq M$, then there exists a saturated chain $M_0(= M)M_1 \cdots M_k(= M')$ in $\mathbf{M}(G)$ between M and M' . Then M_{i-1} covers M_i , and $f_i := M_{i-1} \oplus M_i$ is a proper M_{i-1} -alternating face, $i = 1, 2, \dots, k$. Then we say that M_i is obtained from M_{i-1} by a *Z-transformation* on the (proper M_{i-1} -alternating) face f_i , or simply by *transforming* f_i . Further, we also say that M' is obtained from M by a *Z-transformation sequence* on inner faces $f_1, f_2, \dots, f_k$, and $f_1, f_2, \dots, f_k$ is a face sequence of G by a Z-transformations sequence from M to M' . The Z-transformation of G is *simple* if every inner face of G is transformed at most once during any Z-transformation sequence of G .

Lemma 5.2. *Let $G \in \mathcal{G}$. If $f' \preceq f$ in $\mathbf{F}(G)$, then f' always appears after f in any Z-transformation sequence of G from M^1 to M^0 . Hence Z-transformation of G is simple.*

Proof. Without loss of generality, suppose that f covers f' in $\mathbf{F}(G)$. That is, (f, f') is an arc of directed inner dual $\vec{G}^\#$. Then (f, f') can be extended to a maximal directed path of $\vec{G}^\#$, which can be further extended to a directed cycle of $\vec{G}^*$, denoted by $\vec{C} := f_0^* e_0^* f_1^* e_1^* \dots f_t^* e_t^* f_0^*$; see Fig. 6. Then $T = \{e_0, e_1, \dots, e_t\}$ is an e-cut of G , each edge e_j is a common edge of f_j and f_{j+1} (subscript module $t+1$), and each e_j is a proper edge of f_{j+1} and improper edge of f_j , $0 \leq j \leq t-1$. For any $M \in \mathbf{M}(G)$, by Lemma 5.1, $|M \cap T| = 1$.

Let $P := M_1 M_2 \cdots M_s$ be a directed path in $\vec{Z}(G)$ from M^1 to M^0 . Then $\delta_P(f) = 1$ for all $f \in \mathcal{F}(G)$ by Lemmas 3.3 and 3.4. Suppose that M_{i+1} is obtained from M_i by a Z-transformation on f_j . It is sufficient to show that $f_j, f_{j+1}, \dots, f_t$ do not appear in Z-transformations from M^1 to M_i . We proceed by induction on j . If $j = 1$, then $e_0 \in M_1, M_2, \dots, M_i$. Hence proper edge e_{k-1} of each f_k , $k \geq 2$, does not belong to $M_1, M_2, \dots, M_i$. Hence, the required holds. By induction hypothesis we have that $f_{j+1}, \dots, f_t$ do not appear in Z-transformations from M^1 to M_{i+1} through M_i . Suppose that $M_{i'+1}$ is obtained from $M_{i'}$ by a Z-transformation on f_{j+1} . Then $i+1 \leq i'$, and proper edge e_j of f_{j+1} belong to all $M_{i+1}, \dots, M_{i'}$. That implies that proper edge e_k of f_{k+1} does not belong to $M_{i+1}, \dots, M_{i'}$ for all $k > j$. Hence $f_{j+1}, \dots, f_t$ do not appear in Z-transformations from M^1 to $M_{i'}$; that is, $f_{j+2}, \dots, f_t$ do not appear in Z-transformations from M^1 to $M_{i'+1}$ through $M_{i'}$, as expected. $\square$

For $G \in \mathcal{G}$, we now define a mapping from $\mathcal{M}(G)$ to $J(\mathbf{F}(G))$. For any $M \in \mathcal{M}(G)$, let




Figure 6. An outerplane bipartite graph with e-cuts (the set of edges intersecting a dashed line).

$\sigma(M)$ denote the set of faces in the face sequence by a Z-transformation sequence from M to $M^{\hat{0}}$. By Lemma 3.4, we have

$$\sigma(M) = \{f \in \mathcal{F}(G) \mid f \text{ is contained in the interior of some cycle in } M \oplus M^{\hat{0}}\}.$$

In particular, $\sigma(M^{\hat{0}}) = \emptyset$, and $\sigma(M^{\hat{1}}) = \mathcal{F}(G)$ since $M^{\hat{1}} \oplus M^{\hat{0}}$ is just the boundary of G by Lemma 3.3.

Lemma 5.3. $\sigma : \mathcal{M}(G) \rightarrow J(\mathbf{F}(G))$ is an injective mapping.

Proof. For $M \in \mathbf{M}(G)$, let $f \in \sigma(M)$. If $f' \prec f$ in $\mathbf{F}(G)$, then by Lemma 5.2, f' always appears after f in any Z-transformation sequence of G from $M^{\hat{1}}$ to $M^{\hat{0}}$ passing through M . So $f' \in \sigma(M)$, and $\sigma(M)$ is an order ideal of $\mathbf{F}(G)$. That is, σ is a mapping from $\mathcal{M}(G)$ to $J(\mathbf{F}(G))$. Further it is clear that σ is injective. $\square$

Further, we will show that σ is an isomorphism between $\mathbf{M}(G)$ and $\mathbf{J}(\mathbf{F}(G))$ in the following theorem.

Theorem 5.4. For each $G \in \mathcal{G}$, $\mathbf{M}(G) \cong \mathbf{J}(\mathbf{F}(G))$.

Proof. We first show that, for each $M \in \mathcal{M}(G)$, if $M_1, M_2, \dots, M_k$ are the 1-factors covered by M , then the order ideals $\sigma(M_1), \sigma(M_2), \dots, \sigma(M_k)$ in $\mathbf{M}(G)$ are exactly the ones covered by $\sigma(M)$. By the fact that the order ideals in a finite poset $\mathbf{P}$ covered in $J(P)$ by an order ideal Y are exactly the sets $Y \setminus \{x\}$ for all maximal elements x of Y , it is sufficient to prove that the faces which can be properly transformed in 1-factor M are exactly the maximal elements of $\sigma(M)$.

Let f_i denote properly M -alternating facial cycle such that $f_i = M \oplus M_i$, $i = 1, 2, \dots, k$. For convenience, we also use f_i to denote the corresponding inner face. Hence $\sigma(M_i) =$

$\sigma(M) \setminus \{f_i\}$. By Lemma 5.2, all faces which greater than f_i must be transformed during any transforming sequence from $M^{\hat{1}}$ to M . So each f_i is an maximal element in $\sigma(M)$, and $\sigma(M_i) = \sigma(M) \setminus \{f_i\}$ is covered by $\sigma(M)$.

If f is a maximal element of $\sigma(M)$, then by Lemma 5.2, all elements which greater than f in $\mathbf{F}(G)$ have been transformed from $M^{\hat{1}}$ to M . Let e be any proper edge of f . Let f' be the face of G that has a common edge e with f . If f' is the outer face of G , by Lemma 3.3 e remains unchanged in any Z-transformation from $M^{\hat{1}}$ to M . Otherwise, $f' \in \mathcal{F}(G)$ covers f , and $f' \in \mathcal{F}(G) \setminus \sigma(M)$. Then $e \in M$ since f' has been transformed but f not from $M^{\hat{1}}$ to M and e is an improper edge of f' . Hence, all proper edges of f belong to M ; that is, f is proper M -alternating, as expected.

Further, σ is surjective since $\mathcal{F}(G) = \sigma(M^{\hat{1}})$ is the maximum element of $J(\mathbf{F}(G))$. Therefore σ is an isomorphism between $\mathbf{M}(G)$ and $\mathbf{J}(\mathbf{F}(G))$. $\square$

Given an undirected tree $T = (V, E)$, $\vec{T}$ is any orientation of T . Of course, directed tree $\vec{T}$ has no (directed) cycles. Similar to $\mathbf{F}(G)$, we could consider $\vec{T}$ as the Hasse diagram of a poset $\mathbf{T}$. As a consequence, we obtain another irreducible MDL described in the following result.

Theorem 5.5. *Let $\mathbf{T}$ be a poset derived from any orientation $\vec{T}$ of a tree T . Then $\mathbf{J}(\mathbf{T})$ is an irreducible MDL.*

Proof. By Theorem 5.4, it is sufficient to show that there is a $G \in \mathcal{G}$ such that $\vec{G}^{\#} \cong \vec{T}$. If $|V(T)| \leq 2$, it is obvious. So let $|V(T)| \geq 3$. Let Δ denote the maximum degree of T . We now construct such a graph G as follows. For any vertex v of T , we gave an inner face f_v bounded by a cycle of length 2Δ . If a vertex u of T is adjacent to v , then we place the corresponding inner face f_u outside f_v by overlapping their edges $e' \in f_u$ and $e'' \in f_v$ to a new edge $e \in G$, satisfying the orientation rule of $\mathbf{F}(G)$: (u, v) is an arc from u to v if and only if e goes from the black end-vertex to the white one along the clockwise orientation of f_u . Since f_v has 2Δ edges and v has at most Δ going-out (going-in) arcs in the directed $\vec{T}$, for all other neighbors of v we can proceed similarly. By repeating the above process, one can construct an outerplane bipartite graph $G \in \mathcal{G}$ such that $\vec{G}^{\#} \cong \vec{T}$. For example, see Fig. 7. $\square$

In fact, in the above construction the face degree of f_v may be smaller than 2Δ . The least value of face degree of f_v may reach $2\max\{\Delta_v^-, \Delta_v^+, 2\}$, where Δ_v^-, Δ_v^+ are in- or out-degree of v in $\vec{T}$. During the process, we may need to exchange the order of in- and




Figure 7. Two outerplane bipartite graphs with the orientation for their inner duals.

out-edges such that the edges surrounded f_v are alternately in- and out-edges as many as possible; See Fig. 7 (b).

References

- [1] G. Birkhoff, Lattice Theory, 3rd ed., American Math. Soc., Providence, 1979.
- [2] S.J. Cyvin, I. Gutman, Kekulé Structures in Benzenoid Hydrocarbons, Lecture Notes in Chemistry 46, Springer-Verlag, Berlin, 1988.
- [3] S.J. Cyvin, I. Gutman, Topological properties of benzenoid systems - part XXXVI - Algorithm for the number of Kekulé structures in some peri-condensed benzenoids, MATCH-Commun. Math. Comput. Chem. 19 (1986) 229.
- [4] S. Felsner, Lattice structures from planar graphs, Elec. J. Combin. 11 (2004) #R15.
- [5] J.C. Fournier, Combinatorics of perfect matchings in plane bipartite graphs and application to tilings, Theoret. Comput. Sci. 303 (2003) 333–351.
- [6] G. Grätzer, Lattice Theory, First Concepts and Distributive Lattice, W. H. Freeman and Company, San Francisco, 1971.
- [7] S. Khuller, J. Naor, P. Klein, The lattice structure of flow in planar graphs, SIAM J. Discrete Math. 6 (1993) 477–490.
- [8] S. Klavžar, P. Žigert, A min-max result on catacondensed benzenoid graphs, Appl. Math. Lett. 15 (2002) 279–283.
- [9] S. Klavžar, P. Žigert and G. Brinkmann, Resonance graphs of catacondensed even ring systems are median, Discrete Math. 253 (2002) 35–43.
- [10] P.C.B. Lam and H. Zhang, A distributive lattice on the set of perfect matchings of a plane bipartite graph, Order 20 (2003) 13–29.

- [11] L. Lovász, M. D. Plummer, Matching Theory, Annals of Discrete Mathematics, Vol. 29, North-Holland, Amsterdam, 1986.
- [12] M. Latapy, C. Magnien, Coding distributive lattices with edge firing games, Inf. Process. Lett. 83 (2002) 125–128.
- [13] O. Pretzel, On reorienting graphs by pushing down maximal vertices—II, Discrete Math. 270 (2003) 227–240.
- [14] J. Propp, Lattice structure for orientations of graphs, Manuscript 1993. <http://www.math.wisc.edu/~propp/orient.html>.
- [15] M. Randić, Aromaticity of polycyclic conjugated hydrocarbons, Chem. Rev. 103 (2003) 3449–3605.
- [16] M. Randić, Resonance in catacondensed benzenoid hydrocarbons, Int. J. Quantum Chem. 63(2) (1997) 585–600.
- [17] E. Rémila, The lattice structure of the set of domino tilings of a polygon, Theoret. Comput. Sci. 322 (2004) 409–422.
- [18] E. Rémila, On the structure of some spaces of tilings, SIAM J. Discrete Math. 16 (2002) 1–19.
- [19] N.C. Saldanha, C. Tomei, M.A. Casarin Jr., D. Romualdo, Space of domino tilings, Discrete Comput. Geom. 14 (1995) 207–233.
- [20] W.C. Shiu, P.C.B. Lam, F. Zhang and H. Zhang, Normal components, Kekulé patterns and Clar patterns in plane bipartite graphs, J. Math. Chem. 31 (2002) 405–420.
- [21] P.R. Stanley, Enumerative Combinatorics, Vol. 1, Wadsworth, Belmont, CA, 1986.
- [22] R.J. Wilson, Introduction to Graph Theory, 3rd ed., Longman, New York, 1995.
- [23] F. Zhang, X. Guo and R. Chen, Z -transformation graph of perfect matching of hexagonal systems, Discrete Math. 72 (1988) 405–415.
- [24] F. Zhang, X. Guo and R. Chen, The connectivity of Z -transformation graph of perfect matching of hexagonal systems, Acta Math. Appl. Sinica. 4(2) (1988) 131–135.
- [25] F. Zhang, X. Li, Hexagonal systems with forcing edges, Discrete Math. 140 (1995) 253–263.
- [26] F. Zhang and H. Zhang, A new enumeration method for Kekulé structures of hexagonal system with forcing edges, J. Mol. Struct. (Theochem) 331 (1995) 255–260.
- [27] H. Zhang, The connectivity of Z -transformation graph of perfect matchings of polyominoes, Discrete Math. 158 (1996) 257–272.

- [28] H. Zhang, Direct sum of distributive lattices on the perfect matchings of a plane bipartite graph, *Order* 27 (2) (2010) 101-113.
- [29] H. Zhang, Z-transformation graphs of perfect matchings of plane bipartite graphs: a survey, *MATCH Commun. Math. Comput. Chem.* 56 (2006) 457-476.
- [30] H. Zhang, P.C.B. Lam and W.C. Shiu, Resonance graphs and a binary coding for the 1-factors of benzenoid systems, *SIAM J. Discrete Math.* 22 (2008) 971-984.
- [31] H. Zhang, H. Yao, D. Yang, A min-max result on outerplane bipartite graphs, *Appl. Math. Lett.* 20(2) (2007) 199-205.
- [32] H. Zhang and F. Zhang, The rotation graphs of perfect matchings of plane bipartite graphs, *Discrete Appl. Math.* 73 (1997) 5-12.
- [33] H. Zhang and F. Zhang, Block graphs of Z-transformation graphs of perfect matchings of plane elementary bipartite graphs, *Ars Combin.* 53 (1999) 309-314.
- [34] H. Zhang and F. Zhang, Plane elementary bipartite graphs, *Discrete Appl. Math.* 105 (2000) 291-311.
- [35] H. Zhang, F. Zhang and H. Yao, Z-transformation graphs of perfect matchings of plane bipartite graphs, *Discrete Math.* 276 (2004) 393-404.